

iMedix: Your Personal Health Advisor.

Nulytely Lemon-Lime 420 Gram Oral Solution Hypertonic Laxatives

Search for medical details concerning Nulytely Lemon-Lime oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Nausea, bloating, or feelings of fullness in the stomach/abdomen are very common. Stomach/abdominal cramps, vomiting, and anal irritation occur less often. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: stomach/abdominal pain that is severe or doesn't go away, vomit that is bloody or looks like coffee grounds, chest pain, fainting, fast/irregular heartbeat, sudden shortness of breath, bloody stools, rectal bleeding, seizure, signs of kidney problems (such as change in the amount of urine). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses PEG (polyethylene glycol) with electrolyte is used to clean out the intestines before certain bowel exam procedures such as colonoscopy or barium enema X-rays. It is a laxative that works by drawing large amounts of water into the colon. This effect results in watery bowel movements. Clearing stool from the intestines helps your doctor to better examine the intestines during your procedure. How to use Nulytely Lemon-Lime 420 Gram Oral Solution Hypertonic Laxatives Some brands of this product come with a Medication Guide. If available, read the Medication Guide provided by your pharmacist before you start taking this product. If you have any questions, ask your doctor or pharmacist. Follow any special diet instructions provided by your doctor. Do not eat solid food, usually starting on the day before your procedure until after your procedure. This product must be mixed with water as directed before use. Carefully read and follow all directions to mix this product. After mixing, place the container in the refrigerator. Chilling the mixed liquid will improve its taste. This product may cause certain liquid thickeners (starch-based products) to not work well, which may increase the risk of choking. If you are using liquid thickeners with this product, talk to your pharmacist to select a liquid thickener which will not interact with this product. The dosage is based on your medical condition, procedure, and response to treatment. Drink the mixture by mouth as

directed by your doctor. It is best to drink the mixed liquid quickly instead of sipping slowly. The mixture may also be given through a nose-to-stomach (nasogastric) tube. Follow the specific directions given by your doctor for when and how much to drink. Be sure to finish drinking the amount directed by your doctor. Watery bowel movements usually begin within 1 hour after you start to drink the mixed liquid. Drink plenty of clear liquids as directed by your doctor to prevent becoming dehydrated. Avoid taking other medications by mouth within 1 hour before you start taking this product. Doing so may make the other medication work less well. Ask your doctor or pharmacist if you are not sure when to take your other medications. Precautions Before taking electrolyte with PEG, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: certain stomach/intestinal problems (such as toxic colitis, obstruction, ulcerative colitis), trouble swallowing (such as poor gag reflex, aspiration), history of vomiting easily/often, heart problems (such as heart failure, irregular heartbeat, QT prolongation in the EKG), kidney problems, seizure disorder, mineral imbalances (such as low level of sodium in the blood). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Children may be more sensitive to the side effects of this drug, especially dehydration. Room temperature solution is better for infants. Children younger than 2 years may be more likely to develop low blood sugar from not eating. Get medical help right away if you notice symptoms of low blood sugar in your child, such as shakiness, unusual sleepiness, abnormal/prolonged crying. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Not applicable. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
